

## Ordering Information for Hypohidrotic Ectodermal Dysplasia with Immunodeficiency (HED-ID) Testing

### Indications for Testing

- Recurrent bacterial, viral, and/or mycobacterial infections, usually accompanied by ectodermal dysplasia, in male infants and young children
- Impaired generation of specific antibodies in response to polysaccharide antigens
- Reduced NK cell cytotoxicity
- Male-limited family history of ED and recurrent, severe, or atypical infections, or a family history of *IKBKG*-related HED-ID

### Ordering Information

Gene	CPT Codes	Test Code
<i>IKBKG</i> ( <i>NEMO</i> )	83891(1) 83892(1) 83898(9) 83904(18) 83909(18) 83912(1)	100601

### Family Testing (single amplicon)

<i>IKBKG</i> ( <i>NEMO</i> )	83891(1) 83892(1) 83898(1) 83904(2) 83909(2) 83912(1)	100601
------------------------------	---	--------

### Test Methodology

- Amplification by polymerase chain reaction (PCR); sequencing of entire protein-coding region

### Sample Requirements

- For blood samples:
  - 2 mL whole blood in EDTA tube (lavender top)
  - Samples can be stored briefly at 4°C, but should be shipped on day of collection.
- For buccal swab samples:
  - Please contact client services at 1-866-647-0735 for instructions.
- All sample types should be shipped overnight at room temperature.
- To request a sample shipping kit, please call 1-866-647-0735.

### Turn-around Times

Turn-around times typically range from 7 to 21 days of receipt of sample and all required forms, but may vary depending on test volume and test-specific technical difficulties. Current TATs are posted on our website. Please schedule patient follow-up appointments for discussion of test results conservatively at 6 weeks.

**For more information, please contact Correlagen Diagnostics, Inc., at 1-866-647-0735 or visit us on the web at [www.correlagen.com](http://www.correlagen.com).**